

The equations of caustics for plate problems in bending by using complex potentials

Pericles S. Theocaris and Nikolaos I. Ioakimidis

*Department of Theoretical and Applied Mechanics,
The National Technical University of Athens, Greece*

Abstract The method of caustics is an interesting and efficient experimental method in fracture mechanics and in many more elasticity problems. In plane isotropic elasticity, the equations of the initial curve of the caustic on the specimen and of the caustic itself on the screen originally derived in real forms (the first of them with the help of a Jacobian determinant) were next derived also in complex forms by using the complex potential of Westergaard or, equivalently, the first complex potential of Kolosov–Muskhelishvili. Here this approach is generalized to plate problems in bending now by using the related two complex potentials. The present results are applied to the derivation of the equations of the initial curve of the caustic and of the caustic itself in the problem of bending of an equilateral triangular isotropic elastic plate. Some related remarks are also made.

Final publication details² This technical report was published in 1979 (with quite minor changes) in *Teoretichna i Prilozhna Mekhanika (Journal of Theoretical and Applied Mechanics)*, Sofia. Web address (URL) of the homepage of the *Journal of Theoretical and Applied Mechanics* at De Gruyter: <https://www.degruyter.com/view/j/jtam>. The final publication details of this technical report are

- Theocaris, P. S. and Ioakimidis, N. I., The equations of caustics for plate problems in bending by using complex potentials. *Teoretichna i Prilozhna Mekhanika (Journal of Theoretical and Applied Mechanics)*, Vol. **10** (Year **X**), No. 2, pp. 48–51 (1979).

Therefore, whenever reference to this technical report is to be made, this should be made directly to its above final publication in *Teoretichna i Prilozhna Mekhanika (Journal of Theoretical and Applied Mechanics)*.

1. Introduction

The method of caustics was developed by Theocaris [1] initially for the estimation of stress intensity factors at crack tips in plane isotropic elastic media under generalized plane stress conditions. The equation of the initial curve of the caustic on the specimen was at first determined

¹Both the internal and the external links (all appearing in blue) were added by the second author on 18 March 2018 for the online publication of this technical report.

²These details were also added by the second author on 18 March 2018 for the online publication of this technical report.

by zeroing an appropriate Jacobian determinant J [1] expressed in real form. Also the equation of the caustic itself was initially expressed in real form. In Reference [2], Theocaris proposed the expression of these equations by using the complex potential $Z(z)$ of Westergaard so popular in plane elasticity. This complex potential is equal to the double of the complex potential $\Phi(z)$ of Muskhelishvili. The expression of the equations of caustics through the use of these complex potentials permitted the direct and free from undue calculations application of the method of caustics to a series of engineering problems, most of which are reviewed in Reference [3].

On the other hand, the method of caustics was recently applied to isotropic elastic thin plate problems [4–6]. In such problems, the classical or the Reissner bending theories (the second being more accurate than the first) are generally used [7–9]. In both these methods, the deflection $w(x, y)$ of the points of the plate can be expressed in terms of two complex potentials $\phi(z)$ and $\chi(z)$ exactly as happened with the Airy stress function in plane elasticity [7–9]. In this technical report, we will express the equations of the initial curve of the caustic on the plate and of the caustic itself on the screen in terms of these complex potentials hoping that this will lead to an easier application of the method of caustics to bent plate problems. For simplicity, the plate will be assumed under pure bending conditions, free from any normal loading distribution.

2. Derivation of the equations

In accordance with the developments of References [4, 5], the correspondence between the points (x, y) of the plane of a thin isotropic elastic plate and the corresponding points (u, v) of the screen, when illuminating the plate by a light beam, is established by [4, 5]

$$u = \lambda \left(x + C \frac{\partial w}{\partial x} \right), \quad v = \lambda \left(y + C \frac{\partial w}{\partial y} \right), \quad (1)$$

where $w = w(x, y)$ denotes the deflection of the points of the plate due to bending, λ is the magnification ratio of the optical set-up and C is the overall optical constant of the experimental set-up. Furthermore, in order that a caustic can be formed on the screen, it is necessary that [4, 5]

$$J := \frac{\partial(u, v)}{\partial(x, y)} := \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix} = 0, \quad (2)$$

where J is the related Jacobian determinant. By taking into account Eqs. (1), Eq. (2) can be also written as

$$\begin{vmatrix} 1 + C \frac{\partial^2 w}{\partial x^2} & C \frac{\partial^2 w}{\partial x \partial y} \\ C \frac{\partial^2 w}{\partial x \partial y} & 1 + C \frac{\partial^2 w}{\partial y^2} \end{vmatrix} = 0. \quad (3)$$

The points (x, y) of the plate for which Eq. (3) is satisfied (if such points exist) form the initial curve of the caustic on the plate. The corresponding points on the screen form the caustic itself and are easily determined by using Eqs. (1).

Now, we take into account the fact that both in the classical theory and in the more advanced Reissner theory the deflection $w(x, y)$ of the points of the plate can be expressed (in the absence of normal loading of the plate) in terms of two complex potentials $\phi(z)$ and $\chi(z)$ [7–9], where $z = x + iy$, that is [9, p. 306]

$$w(x, y) = \operatorname{Re} [\bar{z} \phi(z) + \chi(z)] = \frac{1}{2} [\bar{z} \phi(z) + z \overline{\phi(z)} + \chi(z) + \overline{\chi(z)}]. \quad (4)$$

By taking further into account that

$$z = x + iy, \quad \bar{z} = x - iy, \quad \frac{\partial}{\partial x} = \frac{\partial}{\partial z} + \frac{\partial}{\partial \bar{z}}, \quad \frac{\partial}{\partial y} = i \left(\frac{\partial}{\partial z} - \frac{\partial}{\partial \bar{z}} \right), \quad (5)$$

we find that

$$2 \frac{\partial w}{\partial z} = \frac{\partial w}{\partial x} - i \frac{\partial w}{\partial y} = \bar{z} \phi'(z) + \overline{\phi(z)} + \chi'(z). \quad (6)$$

Thus Eqs. (1) can also be written in complex form as

$$u + iv = \lambda \{ z + C [\bar{z} \phi'(z) + \overline{\phi(z)} + \chi'(z)] \}. \quad (7)$$

This equation will determine the form of the caustic on the screen after the points $z = x + iy$ of its initial curve on the plate have been determined.

Hence, we have to take into account Eq. (3) as well. By using Eqs. (4) and (5), we can easily find after some algebra that the equation of the initial curve of the caustic (3) on the plate can be written in complex form as

$$|1 + 2C \operatorname{Re} \phi'(z)| = |C| |\bar{z} \phi''(z) + \chi''(z)|. \quad (8)$$

Equations (8) and (7) can completely determine the shape of the caustic if such a curve is formed on the screen. Evidently, in several cases, Eq. (8) may have no roots.

3. An application

As an application of Eqs. (8) and (7), we consider the problem of an equilateral triangular isotropic elastic plate of height $3c$ and flexural rigidity D loaded by bending moments of intensity M (with $M > 0$) uniformly distributed along its boundary. The deflection $w(x, y)$ in this case is given by [9]

$$w(x, y) = \frac{M}{12cD} [x^3 - 3xy^2 - 3c(x^2 + y^2) + 4c^3], \quad M > 0. \quad (9)$$

By inserting this expression into Eqs. (1) and (2), we find

$$u = \lambda \left[x + \frac{CM}{4cD} (x^2 - 2cx - y^2) \right], \quad v = \lambda \left[y - \frac{CM}{2cD} (x + c)y \right] \quad (10)$$

and

$$\left| 1 - \frac{CM}{2D} \right| = \frac{|C|M}{2D} \frac{r}{c}, \quad r = \sqrt{x^2 + y^2}. \quad (11)$$

From Eq. (11) it is clear that the initial curve of the caustic on the plate is a circle of radius

$$r = c \left| \frac{2D}{CM} - 1 \right|. \quad (12)$$

The same results can be obtained more easily if Eqs. (8) and (7) are used together with the expressions

$$\phi(z) = -\frac{Mz}{4D}, \quad \chi(z) = \frac{M}{12D} \left(\frac{z^3}{c} + 4c^2 \right) \quad (13)$$

for the complex potentials $\phi(z)$ and $\chi(z)$ [9].

4. Some remarks

The use of complex potentials for the solution of bent plate problems is well-known long ago [7–9]. Besides the theoretical results contained in these references, a series of stress concentration problems around holes, where caustics are generally created, have been treated by this method (by assuming the validity of the classical theory of bending of thin plates) in Reference [9]. Recently, Tamate [10] considered crack problems in the theory of bending of thin plates on the basis of the theory of Reissner and using complex potentials. The results of Tamate [10] are interesting not only because the method of caustics is particularly appropriate for the study of crack problems, but also because he formulated these problems through complex functions by using the concept of dislocation distributions. Such a formulation can be proved quite effective for the treatment of almost all general plate bending problems similarly to what has happened in plane elasticity problems. The method of caustics seems to be one of the most powerful experimental techniques to be used together with theoretical results, e.g. for the estimation of stress intensity or stress concentration factors.

References³

- [1] Theocaris, P. S., Local yielding around a crack tip in plexiglas. *Journal of Applied Mechanics*, **37** (2) [*Transactions of the ASME, Series E*, **92** (2)], 409–415 (1970). <https://doi.org/10.1115/1.3408521>
- [2] Theocaris, P. S., Constrained zones in a periodic array of collinear equal cracks. *International Journal of Mechanical Sciences*, **14** (2), 79–94 (1972). [https://doi.org/10.1016/0020-7403\(72\)90090-2](https://doi.org/10.1016/0020-7403(72)90090-2)
- [3] Theocaris, P. S., The topography of entirely or locally curved surfaces studied by caustics. Scientific Publication of the National Technical University of Athens, Athens, 1977.
- [4] Theocaris, P. S. and Gdoutos, E. E., Experimental solution of flexed plates by the method of caustics. *Journal of Applied Mechanics*, **44** (1) [*Transactions of the ASME, Series E*, **99** (1)], 107–111 (1977). <https://doi.org/10.1115/1.3423972>
- [5] Theocaris, P. S., Caustics created from simply supported plates under uniform loading. *International Journal of Solids and Structures*, **13** (12), 1281–1291 (1977). [https://doi.org/10.1016/0020-7683\(77\)90101-9](https://doi.org/10.1016/0020-7683(77)90101-9)
- [6] Theocaris, P. S., Symmetric bending of cracked plates studied by caustics. *International Journal of Mechanical Sciences*, **21** (11), 659–670 (1979). [https://doi.org/10.1016/0020-7403\(79\)90045-6](https://doi.org/10.1016/0020-7403(79)90045-6)
- [7] Green, A. E. and Zerna, W., *Theoretical Elasticity*, 2nd edition. Clarendon Press – Oxford University Press, Oxford, 1968. Web page of the Dover edition: <http://store.doverpublications.com/0486670767.html>
- [8] Solomon, L., *Élasticité Linéaire*. Masson et Cie, Paris. 1968.
- [9] Savin, G. N., *Stress Concentration Around Holes*, English edition (translation of the Russian edition: Gostekhizdat, Moscow – Leningrad, 1951). Pergamon Press, New York – Oxford, 1961. Related web page: https://archive.org/details/nasa_techdoc_19710000647
- [10] Tamate, O., Two arbitrarily situated cracks in an elastic plate under flexure. *International Journal of Solids and Structures*, **12** (4), 287–298 (1976). [https://doi.org/10.1016/0020-7683\(76\)90078-0](https://doi.org/10.1016/0020-7683(76)90078-0)

³All the links (external links in blue) in this section were added by the second author on 18 March 2018 for the online publication of this technical report. Moreover, final publication details were added in Reference [6]